

LINEAR MAPPINGS ON STRONGLY MAGIC SQUARES

Neeradha. C. K.

Dept. of Science & Humanities, Mar Baselios College of Engineering & Technology,
Thiruvananthapuram, Kerala, India, ckneeradha@yahoo.co.in

Abstract: A magic square is a square array of numbers where the rows, columns, diagonals and co-diagonals add up to the same number. Several studies on computational aspects of magic squares are being carried out recently revealing patterns, some of which have lead to analytic insights, theorems or combinatorial results. Magic squares can be used for solving certain complicated and complex problems connected with the algebra and combinatorial geometry of polyhedra, polytopes...While magic squares are recreational on one hand they can be treated somewhat more seriously in higher mathematics on the other hand.

This paper discuss about a well-known class of magic squares; the strongly magic square. The strongly magic square is a magic square with a stronger property that the sum of the entries of the sub-squares taken without any gaps between the rows or columns is also the magic constant. In this paper a generic definition for Strongly Magic Squares is given. The main objective of the paper is to define a function on strongly magic squares which can be established as a linear transformation. The transition of a set of strongly magic squares to an abelian group can be seen in the paper. The paper deals with the formation of a vector space for the set of all strongly magic squares. The kernel of the mapping is obtained as the matrix $\left[A - \frac{\rho}{n^2}U\right]$ where $U = 1; \forall i, j = 1, 2 \dots n^2$.

Keywords: Magic Square, Strongly Magic Square, Linear Transformation, Kernel.

2010 Mathematics Subject Classification Code:20K01, 15A035, 15A04, 47B34.

1. Introduction: Magic squares generally fall into the realm of recreational mathematics (Pasles, 2008), (Pickover, 2002) however a few times in the past century and more recently, they have become the interest of more-serious mathematicians. Magic squares have spelt fascination to mankind throughout history and all across the globe. A normal magic square is a square array of consecutive numbers from $1 \dots n^2$ where the rows, columns, diagonals and co-diagonals add up to the same number. The constant sum is called magic constant or magic number. Along with the conditions of normal magic squares, strongly magic square of order n have a stronger property that the sum of the entries of the sub-squares taken without any gaps between the rows or columns is also the magic constant. The study on numerical properties of strongly magic squares of order n have been carried out by astrologer turned mathematician Padmakumar (Padmakumar, 1995). Another study carried out by Stanley (Richard P. Stanley, 1978) on magic Squares using the tools of Commutative Algebra which makes use of graded rings to define a hilbert series (Qimh Richey Xantcha, 2012). In this paper some advanced mathematical properties of the strongly magic squares are discussed.

2. Notations and Mathematical Preliminaries

2.1) **Magic Square:** A magic square of order n over a field R where R denotes the set of all real numbers is an n^{th} order matrix $[a_{ij}]$ with entries in R such that

$$\sum_{j=1}^n a_{ij} = \rho \quad \text{for } i = 1, 2, \dots, n \quad (1)$$

$$\sum_{j=1}^n a_{ji} = \rho \quad \text{for } i = 1, 2, \dots, n \quad (2)$$

$$\sum_{i=1}^n a_{ii} = \rho, \quad \sum_{i=1}^n a_{i,n-i+1} = \rho \tag{3}$$

Equation (1) represents the row sum, equation (2) represents the column sum, equation (3) represents the diagonal and co-diagonal sum and symbol ρ represents the magic constant. (Small, 1988)

2.2) **Magic Constant:** The constant ρ in the above definition is known as the magic constant or magic number. The magic constant of the magic square A is denoted as $\rho(A)$. In the example given below the magic constant of A is 15 and B is 34

$$A = \begin{array}{|c|c|c|} \hline 8 & 1 & 6 \\ \hline 3 & 5 & 7 \\ \hline 4 & 9 & 2 \\ \hline \end{array}$$

$$B = \begin{array}{|c|c|c|c|} \hline 9 & 16 & 5 & 4 \\ \hline 7 & 2 & 11 & 14 \\ \hline 12 & 13 & 8 & 1 \\ \hline 6 & 3 & 10 & 15 \\ \hline \end{array}$$

2.3) **Strongly magic square (SMS): Generic Definition:** A strongly magic square over a field R is a matrix $[a_{ij}]$ of order $n^2 \times n^2$ with entries in R such that

$$\sum_{j=1}^{n^2} a_{ij} = \rho \text{ for } i = 1, 2, \dots, n^2 \tag{4}$$

$$\sum_{i=1}^{n^2} a_{ji} = \rho \text{ for } i = 1, 2, \dots, n^2 \tag{5}$$

$$\sum_{i=1}^{n^2} a_{ii} = \rho, \quad \sum_{i=1}^{n^2} a_{i,n^2-i+1} = \rho \tag{6}$$

$$\sum_{l=0}^{n-1} \sum_{k=0}^{n-1} a_{i+k, j+l} = \rho \text{ for } i, j = 1, 2, \dots, n^2 \tag{7}$$

where the subscripts are congruent modulo n^2

Equation (4) represents the row sum, equation (5) represents the column sum, equation (6) represents the diagonal & co-diagonal sum, equation (7) represents the $n \times n$ sub-square sum with no gaps in between the elements of rows or columns and is denoted as $M_{OC}^{(n)}$ or $M_{OR}^{(n)}$ and ρ is the magic constant.

Note: The n^{th} order subsquare sum with k column gaps or k row gaps is generally denoted as $M_{kC}^{(n)}$ or $M_{kR}^{(n)}$ respectively (Neeradha.C.K, 2016)

2.4) **Kernel of A Homomorphism:** If ϕ is a homomorphism of a group G into G' , then the kernel of ϕ is denoted as $\ker \phi$ and is defined as $\ker \phi = \{g \in G; \phi(g) = e'\}$, where e' is the identity of G'

2.5) **Linear Transformation:** Let U and V be two vector spaces over the same field F . Then a mapping $f: U \rightarrow V$ is called linear transformation of U into V if $f(\lambda a + \mu b) = \lambda f(a) + \mu f(b) \quad \forall \lambda, \mu \in F$ and $a, b \in U$ (Kenneth Hoffmann, 1971)

2.6) **Other Notations:**

1. R denotes the set of all real numbers.
2. S denote the set of all strongly magic squares of order $n^2 \times n^2$

3. Propositions and Theorems:

Proposition 3.1: If A and B are two Strongly magic squares of order $n^2 \times n^2$ with $\rho(A) = a$ and $\rho(B) = b$, then $C = (\lambda + \mu)(A + B)$ is also a Strongly magic square with magic constant $(\lambda + \mu)(\rho(A) + \rho(B))$; for every $\lambda, \mu \in R$

Proof: Let $A = [a_{ij}]_{n^2 \times n^2}$ and $B = [b_{ij}]_{n^2 \times n^2}$

Then $C = (\lambda + \mu)(A + B)$

$$= [(\lambda + \mu)(a_{ij} + b_{ij})]$$

Sum of the i^{th} row elements of

$$C = \sum_{j=1}^{n^2} c_{ij} = (\lambda + \mu) \left(\sum_{j=1}^{n^2} (a_{ij}) + \sum_{j=1}^{n^2} (b_{ij}) \right)$$

$$= (\lambda + \mu)(a + b)$$

$$= (\lambda + \mu)(\rho(A) + \rho(B))$$

A similar computation holds for column sum, diagonals sum and sum of the $n \times n$ sub squares

From the above propositions the following results can be obtained by putting suitable values for λ , and μ

Results:

If for every $\lambda, \mu \in R$ and $A, B \in S$,

1.1) $\lambda(A + B) \in S$ with $\rho(\lambda(A + B)) = \lambda(\rho(A) + \rho(B))$

1.2) $(A + B) \in S$ with $\rho(A + B) = \rho(A) + \rho(B)$

1.3) $\lambda A \in S$ with $\rho(\lambda A) = \lambda \rho(A)$

1.4) $\lambda A + \mu B \in S$ with $\rho(\lambda A + \mu B) = \lambda \rho(A) + \mu \rho(B)$

1.5) $-A \in S$ with $-A \in S$

Theorem 3.2: $\langle S, + \rangle$ forms an abelian group. (MadhukarMallayya, Neeradha. C. K , 2016)

Proof:

a) Closure property: if $A, B \in S$, then $A + B \in S$. (from above result 1.2)

b) Associativity : if $A, B, C \in S$, then $A + (B + C) = (A + B) + C \in S$ (Since matrix addition is associative.)

c) Existence of Identity: There exists o matrix in S so that $A + 0 = 0 + A = A$, where 0 acts as the identity element.

d) Existence of additive inverse : For every $A \in S$, there exists $-A \in S$ so that $A + (-A) = 0$ where $0 \in S$ (from result 1.5).

e) Commutativity :If $A, B \in S$, then $A + B = B + A \in S$ (Since matrix addition is commutative .)

This completes the proof.

Proposition 3.3: For all $A, B \in S, \lambda, \mu \in R$;

a) $\lambda(A + B) = \lambda A + \lambda B$

b) $(\lambda + \mu).A = \lambda.A + \mu.A$

c) $(\lambda\mu).A = \lambda.(\mu.A)$

d) $1.A = A$

Proof: Since $A, B \in S; A = [a_{ij}]_{n^2 \times n^2}$ and $B = [b_{ij}]_{n^2 \times n^2}$

1. $A + B = [a_{ij} + b_{ij}]$

$$\lambda(A + B) = \lambda[a_{ij} + b_{ij}]$$

$$= [\lambda a_{ij} + \lambda b_{ij}]$$

$$= [\lambda a_{ij}] + [\lambda b_{ij}]$$

$$= \lambda [a_{ij}] + \lambda [b_{ij}]$$

$$= \lambda . A + \lambda . B$$

Similarly results (ii),(iii), and (iv) can be proved.

Theorem 3.4: $\langle S, +, . \rangle$ forms a vector space over the field of real numbers

Proof: It is an immediate consequence of Theorem.3.2 and Proposition 3.3 (Neeradha.C.K,V.Madhukar.Mallayya, 2016)

Proposition 3.5: The mapping $\phi : S \rightarrow R$ defined by $\phi(A) = \rho(A)$, $\forall A \in S$ is a linear transformation

Proof: Let $A, B \in S$

$$\begin{aligned} \phi(\lambda A + \mu B) &= \rho(\lambda A + \mu B) = \lambda \rho(A) + \mu \rho(B) \quad (\text{By Result 1.4 and Theorem 3.4}) \\ &= \lambda \phi(A) + \mu \phi(B) \end{aligned}$$

Proposition 3.6: The kernel of the mapping $\phi : S \rightarrow R$ defined by $\phi(A) = \rho(A)$, $\forall A = [a_{ij}] \forall i, j = 1, 2 \dots n^2 \in S$ is $\text{Ker } \phi = A' = \left[A - \frac{\rho}{n^2} U \right]$ where $\rho(A) = \rho$ and $U = [u_{ij}]$ such that $u_{ij} = 1 \forall i, j = 1, 2 \dots n^2$

Proof: Let $A' = \left[A - \frac{\rho}{n^2} U \right] = \left[a_{ij} - \frac{\rho}{n^2} u_{ij} \right]$

$$\begin{aligned} \text{Ker } \phi &= \left\{ A' \in S \text{ such that } \phi(A') = 0 \right. \\ &= \left. \{ A' \in S \text{ such that } \rho(A') = 0 \right. \end{aligned}$$

Now

$$\rho(A') = \sum_{j=1}^{n^2} a_{ij} - \rho = 0$$

$$\text{Therefore } A' = \left[A - \frac{\rho}{n^2} U \right] \subset \text{Ker } \phi$$

Now let $B = [b_{ij}] \forall i, j = 1, 2 \dots n^2 \in \text{Ker } \phi$, then $\rho(B) = 0$

$$\text{Clearly } B = [b_{ij}] = \left[b_{ij} - \frac{0}{n^2} u_{ij} \right] \subset A'$$

$$\text{Therefore } \text{Ker } \phi = A' = \left[A - \frac{\rho}{n^2} U \right]$$

4. Conclusion: The study of strongly magic squares is an emerging innovative area in which mathematical analysis can be done. Here some advanced properties regarding strongly magic squares namely Abelian group structure, vector spaces, group homomorphism, group isomorphism, vector space isomorphism, linear transformation, kernel of transformation are described. Physical application of magic squares is still a new topic that needs to be explored more. Ollerenshaw and BrËe (Ollerenshaw, 1999) have a patent for using most-perfect magic squares for cryptography, and Besslich (Besslich, 1983), (Besslich, Ph. W, 1983) has proposed using pan diagonal magic squares as dither matrices for image processing. Further studies are being carried out by the authors on the scope for further research and the application of Strongly magic squares on Diophantine equations, Moment of inertia, Electric Quadrapoles, Data hiding Schemes etc.

5. Acknowledgement: We express sincere gratitude for the valuable suggestions given by Dr. Ramaswamy Iyer, Former Professor in Chemistry, Mar Ivanios College, Trivandrum, in preparing this paper.

References:

1. Besslich, P. W. (1983). Comments on electronic techniques for pictorial image reproduction. IEEE Trans. on Comm., 31, 846-847.
2. Besslich, Ph. W. (1983). A Method for the Generation and Processing of Dyadic Indexed Data. IEEE Trans. on Computers, C-32(No. 5), 487-494.
3. Kenneth Hoffmann, K. (1971). Linear Algebra (2nd edition ed.). London: Prentice Hall International, INC.
4. MadhukarMallayya, Neeradha. C. K . (2016, December). Strongly Magic Squares As Abelian Groups. Int. J. of Mathematical Sciences and Applications, Vol. 6 (No. 2), pp 559-563.
5. Neeradha.C.K. (2016). Magic Squares - A Brief Note on its Applications. Bulletin of Kerala Mathematical Association, 209-215.
6. Neeradha.C.K,V.Madhukar.Mallayya. (2016, August). Strongly Magic Square As Vector Space. International J. of Pure & Engg. Mathematics, Vol. 4 (No. II), pp. 1-9.
7. Ollerenshaw, K. a. (1999, June 17). Patent No. WO 99/66671 Patent.
8. Padmakumar, T. (1995). Strongly Magic Square, Applications Of Fibonacci Numbers. Volume 6. Proceedings of The Sixth International Research Conference on Fibonacci Numbers and Their Applications.

9. Pasles, P. C. (2008). Benjamin Franklin's numbers: an unsung mathematical odyssey. Princeton, NJ: Princeton University Press.
10. Pickover, C. (2002). The Zen of Magic Squares, Circles and Stars. Princeton, NJ: Princeton University Press.
11. Qimh Richey Xantcha, J. A. (2012, August). Stanley's Theory of Magic Squares. Retrieved from <http://www2.math.uu.se/~qimh/Magic.pdf>
12. Richard P. Stanley. (1978). Hilbert Functions of Graded Algebras. *Advances in Mathematics* 28, 57-83.
13. Small, C. (1988, Aug. - Sep). Magic Squares Over Fields. *The American Mathematical Monthly*, Vol. 95(No. 7), pp. 621-625.
